

.....

/ / /

(Zea mays L.)

. .

()
)

)

()
(

(

.

.

.

/ /

.

/ /

.

/ /

.

.

:

-

-

.....

(*Zea mays* L.)

(ÿ)

()

ÿ

()

()

/

()

()

ÿÿ

-)

/

(

()

/

ÿ

.....

/ / /

()

/

/

/ ÿ /

()

()

ÿ

()

()

ÿ

ÿ

ÿ

(*Zea mays* L.)

)

)

(K3640/5 A679 K3651/1)

K47/2-2-1-21-2-1-1- K19)

(1

(K18 K166B K166A)

:

()

()

)

(

ÿ

()

)

(

ÿÿ

)

ÿ

(

ÿÿÿ

..... / / /

ÿÿ Dickey John

GAC

.()

()

()

SPSS

K18×K166B

ÿ

K18×K166B

ÿ/

K166B×K3640/5

.()

K18×K166B

K18×K47/2-2-1-21-2-1-1-1

()

() .()

()

()

(%)

()

cdef	defg	bcdef	defg	abcdefg	fgh	yy ^{bcde}	h	K18×K3651/1
cdef	cdefg	bc	cdefg	abcdefg	y ^{bcdef}	y ^{bcde}	gh	K18×A679
cdef	bcdef	cdef	cdefg	ab	y ^{bcdef}	y ^{abcde}	defg	K18×K166A
cdef	a	bcd	a	abcdefg	abcd	y ^a	a	K18×K166B
cdef	bedef	bcdef	cdefg	ab	efg	bcde	defg	K18×K3640/5
cdef	a	y ^{bcdef}	b	ab	a	abc	ab	K18×K47/2-2-1-21-2-1-1-1
defg	bc	bcdef	y ^c	g	abcde	yy ^{ef}	y ^{def}	K18×K19
fgh	defg	fg	defg	abcdefg	efg	abcd	y ^h	K3651/1×A679
efgh	bcdef	efg	cdefg	abcdefg	abcde	y ^{cdef}	efgh	K3651/1×K166A
cde	y ^{bcde}	bc	cde	abcdefg	abcde	yy ^{bcdef}	cde	K3651/1×K166B
a	bcde	a	cde	ab	efg	y ^{abc}	defg	K3651/1×K3640/5
bcd	efg	b	efg	ab	gh	abcde	gh	K3651/1×K47/2-2-1-21-2-1-1-1
cdef	g	bcdef	g	ab	h	abcde	gh	K3651/1×K19
b	bcdef	a	cdefg	abcdefg	y ^{bcdef}	y ^{bcdef}	fgh	A679×K166A
y ^{cdef}	g	defg	g	ab	gh	y ^f	y ^h	A679×K166B
y ^{gh}	fg	y ^g	fg	abcdefg	h	yy ^{abcde}	fgh	A679×K3640/5
fgh	efg	fg	fg	abcdefg	y ^{bcdef}	abcde	defg	A679×K47/2-2-1-21-2-1-1-1
bc	fg	bc	fg	ab	fgh	def	h	A679×K19
cdef	bcd	bc	b	abcdefg	abcde	abc	cde	K166A×K166B
cdef	y ^{bcde}	bedde	cdefg	ab	y ^{bcdef}	ab	defg	K166A×K3640/5
defg	bedef	defg	cdefg	ab	efg	y ^{abc}	defg	K166A×K47/2-2-1-21-2-1-1-1
y ^{cdef}	bc	y ^{cdef}	y ^c	efg	a	abc	cde	K166A×K19
efgh	y ^{bcde}	fg	cde	abcdefg	abcde	abcde	y ^{cde}	K166B×K3640/5
cdef	bdefg	cdefg	cdefg	abcdefg	abcde	yy ^{abc}	cd	K166B×K47/2-2-1-21-2-1-1-1
cdef	b	bcd	y ^c	abcdefg	a	yy ^{bcde}	y ^{bc}	K166B×K19
h	defg	y ^g	defg	efg	y ^{bcdef}	abcde	y ^{defg}	K3640/5×K47/2-2-1-21-2-1-1-1
cde	g	bcd	efg	ab	fgh	bcde	defg	K3640/5×K19
cdef	a	y ^{bcdef}	ab	efg	a	y ^{bcdef}	defg	K47/2-2-1-21-2-1-1-1×K19

		()	()	()						
cd	def	/ abcd	/ fghij	ŷ/ b	ŷ/ f	cde	fgh			K18×K3651/1
cd	def	/ bcde	/ ghij	ŷ/ b	ŷ/ def	cdef	efgh			K18×A679
ŷŷ ^{abcd}	ef	/ ŷe	/ abcde	ŷ/ b	ŷ/ ef	ŷ ^a	abcdef			K18×K166A
a	a	/ e	ŷ ⁱ	ŷ/ b	ŷ/ a	abcd	abcde			K18×K166B
abcd	bcde	/ abcd	/ ab	ŷ/ b	ŷ/ f	cd	ŷ abc			K18×K3640/5
abcd	a	/ ŷabcd	ŷ/ j	ŷ/ b	ŷ/ abcd	cde	abcdef			K18×K47/2-2-1-21-2-1-1-1
ab	bcde	/ cde	ŷ/ ij	ŷ/ b	ŷ/ cdef	bcd	a			K18×K19
ŷ ^d	f	/ ŷ ^{abcd}	/ abcdefg	ŷ/ b	ŷ/ cdef	cde	abcdef			K3651/1×A679
abcd	def	/ ŷ ^{ab}	/ abcdef	ŷ/ b	ŷ/ def	cdef	abcdef			K3651/1×K166A
abcd	cdef	/ abc	/ bcdefg	ŷ/ b	ŷ/ ef	def	ŷ bcdefg			K3651/1×K166B
bcd	def	/ ŷ ^a	/ abcd	ŷ/ b	ŷ/ cdef	def	fgh			K3651/1×K3640/5
ŷ ^{abcd}	def	/ abcd	/ abcd	ŷ/ b	ŷ/ f	ŷ ^{def}	fgh			K3651/1×K47/2-2-1-21-2-1-1-1
abcd	ŷ ^{cdef}	/ ŷ ^{abc}	/ abc	ŷ/ b	ŷ/ f	def	ŷ cdefgh			K3651/1×K19
abcd	def	/ a	/ abcdefg	ŷ/ b	ŷ/ bcde	cde	abcdef			A679×K166A
ŷ ^{abcd}	def	/ abc	/ abcd	ŷ/ b	ŷ/ f	ŷ ^f	h			A679×K166B
abcd	cdef	/ abcd	/ abcdef	ŷ/ b	ŷ/ f	cd	gh			A679×K3640/5
d	ŷ ^{def}	/ de	/ hij	ŷ/ b	ŷ/ ef	ef	gh			A679×K47/2-2-1-21-2-1-1-1
cd	ef	/ abcd	/ a	b	ŷ/ ef	cd	fgh			A679×K19
ŷ ^{abc}	bcd	/ abcd	/ fghij	ŷ/ b	ŷ/ cdef	abc	ŷ abcd			K166A×K166B
abcd	bcde	/ ŷabcd	/ abcdef	ŷ/ b	ŷ/ bcde	cde	fgh			K166A×K3640/5
ŷ ^{abcd}	bcde	/ ab	/ abcdef	ŷ/ b	ŷ/ def	def	ab			K166A×K47/2-2-1-21-2-1-1-1
ŷŷ ^{abcd}	ŷ ^{bcde}	/ abcd	/ defghij	b	ŷ/ ab	ab	abcdef			K166A×K19
abcd	def	e	/ defghij	/ a	ŷ/ cdef	abcd	ŷ abcd			K166B×K3640/5
abc	b	/ abc	/ defghij	ŷ/ b	ŷ/ cdef	abcd	abcdef			K166B×K47/2-2-1-21-2-1-1-1
abcd	ŷ ^{bc}	/ abcd	/ efghij	ŷ/ b	ŷ/ abc	abcd	abcde			K166B×K19
abcd	ef	/ ŷabcd	/ hij	ŷ/ b	ŷ/ def	cdef	ŷ ^{defgh}			K3640/5×K47/2-2-1-21-2-1-1-1
ŷ ^{abcd}	ef	/ abc	/ abc	ŷ/ b	ŷ/ def	def	ŷ ^{fgh}			K3640/5×K19
ŷ ^{abcd}	ŷ ^{bcde}	/ abcd	/ abcdef	ŷ/ b	ŷ/ cdef	ŷ ^{cdef}	abcdef			K47/2-2-1-21-2-1-1-1×K19

()

()

..... / / /

/

yy

()

-

	F	R ²	X ₃	X ₂	X ₁	
/	**	yl	-	-	yl	(x ₁)
/	**	yl	-	yl	yl	(x ₂)

()

yl	yl/y	yl y *	yl
yl y	yl	yl *	yl
-yl/y	yl y	yl *	yl
yl	yl	yl *	yl
yl	yl	yl *	yl
-yl	yl	-yl *	yl y
yl	yl y	yl *	yl
yl	-yl	-yl *	yl
yl *	yl	yl y *	yl
yl y	yl y *	yl	yl y
yl	-yl *	yl y	yl
yl	yl *	yl	yl
-yl/y	yl y *	yl	yl
yl *	yl/y	-yl/yy	yl
/	/	/	
/	/	/	

.*

.....

(*Zea mays* L.)

/ /

/ /

/

()

()

()

/ /

()

() ()

()

()

.....

/ / /

()

()

()

ÿ

/ ÿ /

/ÿ

/

% /

()

/

()

K18×K166B

()

-y/y	y/	y/	y/	y/ *	y/
-y/y	y/	-y/	y/ y	y/ *	y/
y/ y	-y/y	y/	-y/	y/ *	y/
y/ y	y/	-y/y	y/ *	y/y	y/
y/	y/y	-y/	y/ *	y/	y/
y/	y/	-y/y	-y/ *	-y/y	y/
y/y	y/y	y/ *	-y/y	y/ yy	y/
y/ y	y/y	y/ *	y/	y/	y/
-y/	y/	y/ *	y/	y/ y	y/
y/y y	-y/y	-y/ *	y/	y/y	y/ y
-y/ y	-y/	-y/ *	y/ y	y/ *	y/
y/y	y/ y *	y/y	-y/y	y/y	y/
-y/	y/ *	y/	-y/y	y/ y	y/
-y/ y *	y/ yy	y/	y/ y	y/	y/
/	/	/	/y	/	
/	/	/ y	/y	/	

.*

- / / /
1. Bahosh, M., B. Rabiei and H. Abasdokht. 2008. Principals components analysis for morphological traits in grain maize hybrids. The 10th Iranian Crop Science and Breeding Congress Iran, Karaj. 159 pp.
 2. Basafa, M. 2004. Study yield and correlation phenotypical different traits with grain yield in new premature maize (*Zea mays* L.) hybrids. The 8th Crop Production and Breeding Congress Iran, Guilan. 16 pp.
 3. Choukan, R., A.A. Hosseinzadeh, M.R. Ghanadha, A.R. Talei and A. Mohammadi. 2005. Classification of maize inbred lines based on morphological traits. *Plant and Seed*. 21(1): 139-157.
 4. Estakhr, A. and R. Choukan. 2006. Study yield component yield and correlation between their in inner and outer hybrids. *Agric. Sci. J.*, 37(1): 85-91.
 5. Galicia, L., E. Nurit, A. Rosales and N. Palacios-Rojas. 2008. Maize nutrition quality and plant tissue analysis laboratory, CIMMYT. 42 pp.
 6. Johnson, C. 2000. Ag answers: post-pollination period critical to maize yeilds. Agricultural Communication Service, Purdue University. 12 pp.
 7. Jones, R.J., A.J. Roessler and S. Ouattar. 1985. Thermal environment during endosperm cell division in maize: Effects on number of endosperm cells and starch granules. *Crop Sci.*, 25: 830-834.
 8. Khajenouri, A. 1996. Advanced statistic and biometry. University Tehran. 467 pp.
 9. Khodarahmpour, Z., R. Choukan, M.R. Bihamta and E. Majidi. 2009. Multivariate Analysis Maize (*Zea mays* L.) Inbred lines in Heat Stress Condition. 1st Regional Conference on Tropical Crops Production under Environmental Stresses Conditions. 38 pp.
 10. Lauer, J. 2006. Concerns about drought as maize pollination begins. *Wiscosin Crop Manager*. 7 pp.
 11. Mohebi, Z., S. Aharizad and M.R. Shiri. 2008. Grouping of early maturity grain maize hybrids using multivariate statistical methods. The 10th Iranian Crop Science & Breeding Congress. 161 pp.
 12. Shoaeh Hosseini, M., M. Farsi and Khavari S. Khorasani. 2008. Study effects water deficit stress on yield and components yield in some grain maize hybrid with use of path analysis. *Agric. Knowledg J.*, 18(1): 71-85.
 13. Wilhelm, E.P., R.E. Mullen, P.L. Keeling and G.W. Singletary. 1999. Heat stress during grain filling in maize effects on kernel growth and metabolism. *Crop Sci.*, 39: 1733-1741.
 14. Zeinali, H., A. Nasrabadi, H. Hosseinzadeh, R. Choukan and M. Sabokdast, 2005. Factor analysis in grain maize hybrid cultivars. *Agric. Sci. J.*, 36(4): 895-902.

An Evaluation of Some Quantitative Traits in Maize (*Zea mays* L.) Hybrids Under Heat Stress Using Multivariate Analysis

Z. Khodarahmpour¹ and R. Choukan²

Abstract

In order to determine the best index for evaluating maize genotypes, 28 maize hybrids were evaluated in two planting dates, 6 July to coincide heat stress with pollination time and 27 July as normal planting to avoid high temperature during pollination and grain filling period, using a randomized complete block design with three replications, in Shushtar city, in 2008. Results showed that in stepwise regression analysis for heat stress condition, grain dry matter weight and grain depth traits were entered in model but, for normal condition, nothing trait werenot entered in model. Factor analysis, for heat stress and normal conditions indentified three and five independent factors which explained 73.59 and 75.78 percent of all variations, respectively. In heat stress condition, first factor, named yield and yield components and in normal condition named ear morphology, explaining 43.78 and 18.96 percent of total variations, respectively. Second factor for heat stress and third factor for normal condition named grain characteristic, which explained 15.67 and 17.58 percent of total variations, respectively. Third factor in heat stress named ear morphology. Second factor, yield components, fourth factor grain width and grain dry matter weight and fifth factor grain yield named in normal condition. Based on the results, grain dry matter weight and grain depth under heat stress revealed as suitable traits which can discriminate maize genotypes.

Keywords: Factor analysis, Heat stress, Maize, Hybrids

1- Assistant Professor, Islamic Azad University Shushtar Branch

2- Associate Professor, Seed and Plant Improvement and Production Research Institute, Karadj